



Protect Yourself When Handling Sharps

U.S. Department of Labor
Occupational Safety and Health Administration



A needlestick or a cut from a contaminated scalpel can lead to infection from hepatitis B virus (HBV) or human immunodeficiency virus (HIV) which causes AIDS. Although few cases of AIDS have been documented from occupational exposure, approximately 8,700 health care workers each year contract hepatitis B. About 200 will die as a result. The new OSHA standard covering bloodborne pathogens specifies measures to reduce these risks of infection.

PROMPT DISPOSAL

The best way to prevent cuts and sticks is to minimize contact with sharps. That means disposing of them immediately after use. Puncture-resistant containers must be available nearby to hold contaminated sharps-- either for disposal or, for reusable sharps, later decontamination for re-use. When reprocessing contaminated reusable sharps, employees must not reach by hand into the holding container. Contaminated sharps must never be sheared or broken.

Recapping, bending, or removing needles is permissible only if there is no feasible alternative or if required for a specific medical procedure such as blood gas analysis. If recapping, bending, or removal is necessary, workers must use either a mechanical device or a one-handed technique. If recapping is essential--for example, between multiple injections for the same patient--employees must avoid using both hands to recap. Employees might recap with a one-handed "scoop" technique, using the needle itself to pick up the cap, pushing cap and sharp together against a hard surface to ensure a tight fit. Or they might hold the cap with tongs or forceps to place it on the needle.

SHARPS CONTAINERS

Containers for used sharps must be puncture resistant. The sides and the bottom must be leakproof. They must be labeled or color coded red to ensure that everyone knows the contents are hazardous. Containers for disposable sharps must have a lid, and they must be maintained upright to keep liquids and the sharps inside.

Employees must never reach by hand into containers of contaminated sharps. Containers for reusable sharps could be equipped with wire basket line for easy removal during reprocessing, or employees could use tongs or forceps to withdraw the contents. Reusable sharps disposal containers may not be opened, emptied, or cleaned manually.

Containers need to be located as near to as feasible the area of use. In some cases, they may be placed on carts to prevent access to mentally disturbed or pediatric patients. Containers also should be available wherever sharps may be found, such as in laundries. The containers must be replaced routinely and not be overfilled, which can increase the risk of needlesticks or cuts.

HANDLING CONTAINERS

When employees are ready to discard containers they should first close the lids. If there is a chance of leakage from the primary container, the employees should use a secondary container that is closable, labeled, or color coded and leak resistant.

Careful handling of sharps can prevent injury and reduce the risk of infection. By following these work practices, employees can decrease their chances of contracting bloodborne illness.