



Holding the Line on Contamination

U.S. Department of Labor
Occupational Safety and Health Administration



Keeping work areas in a clean and sanitary condition reduces employees' risk of exposure to bloodborne pathogens. Each year about 8,700 health care workers are infected with hepatitis B virus, and 200 die from contracting hepatitis B through their work. The chance of contracting human immunodeficiency virus (HIV), the bloodborne pathogen which causes AIDS, from occupational exposure is small, yet a good housekeeping program can minimize this risk as well.

DECONTAMINATION

Every employer whose employees are exposed to blood or other potentially infectious materials must develop a written schedule for cleaning each area where exposures occur. The methods of decontaminating different surfaces must be specified, determined by the type of surface to be cleaned, the soil present and the tasks or procedures that occur in that area.

For example, different cleaning and decontamination measures would be used for a surgical operatory and a patient room. Similarly, hard surfaced flooring and carpeting require separate cleaning methods. More extensive efforts will be necessary for gross contamination than for minor spattering. Likewise, such varied tasks as laboratory analyses and normal patient care would require different techniques for clean-up.

Employees must decontaminate working surfaces and equipment with an appropriate disinfectant after completing procedures involving exposure to blood. Many laboratory procedures are performed on a continual basis throughout a shift. Except as discussed below, it is not necessary to clean and decontaminate between procedures. However, if the employee leaves the area for a period of time, for a break or lunch, then contaminated work surfaces must be cleaned.

Employees also must clean (1) when surfaces become obviously contaminated; (2) after any spill of blood or other potentially infectious materials; and (3) at the end of the work shift if contamination might have occurred. Thus, employees need not decontaminate the work area after each patient care procedure, but only after those that actually result in contamination.

If surfaces or equipment are draped with protective coverings such as plastic wrap or aluminum foil, these coverings should be removed or replaced if they become obviously contaminated. Reusable receptacles such as bins, pails and cans that are likely to become contaminated must be inspected and decontaminated on a regular basis. If contamination is visible, workers must clean and decontaminate the item immediately, or as soon as feasible.

Should glassware that may be potentially contaminated break, workers need to use mechanical means such as a brush and dustpan or tongs or forceps to pick up the broken glass--never by hand, even when wearing gloves.

Before any equipment is serviced or shipped for repairing or cleaning, it must be decontaminated to the extent possible. The equipment must be labeled, indicating which portions are still contaminated. This enables employees and those who service the equipment to take appropriate precautions to prevent exposure.

REGULATED WASTE

In addition to effective decontamination of work areas, proper handling of regulated waste is essential to prevent unnecessary exposure to blood and other potentially infectious materials. Regulated waste must be handled with great care --i.e., liquid or semi liquid blood and other potentially infectious materials, items caked with these materials, items that would release blood or other potentially infected materials if compressed, pathological or microbiological wastes containing them and contaminate sharps.

Containers used to store regulated waste must be closable and suitable to contain the contents and prevent leakage of fluids. Containers designed for sharps also must be puncture resistant. They must be labeled or color coded to ensure that employees are aware of the potential hazards. Such containers must be closed before removal to prevent the contents from spilling. If the outside of a container becomes contaminated, it must be placed within a second suitable container.

Regulated waste must be disposed of in accordance with applicable state and local laws.

LAUNDRY

Laundry workers must wear gloves and handle contaminated laundry as little as possible, with a minimum of agitation. Contaminated laundry should be bagged or placed in containers at the location where it is used, but not sorted or rinsed there.

Laundry must be transported within the establishment or to outside laundries in labeled or red color-coded bags. If the facility uses Universal Precautions for handling all soiled laundry, then alternate labeling or color coding that can be recognized by the employees may be used. If laundry is wet and it might soak through **laundry** bags, then workers must use bags that prevent leakage to transport it.

RESEARCH FACILITIES

More stringent decontamination requirements apply to research laboratories and production facilities that work with concentrated strains of HIV and HBV.

from OSHA regional offices.